Journal of Immunological Methods, 117 (1989) 296–297
Elsevier

Contents to volume 117

Number 1 – 8 February 1989

Research reports

Production of hybrid hybridomas based on HAT^s-neomycin^r double mutants (JIM 05043)
W.B.M. De Lau, A.E. Van Loon, K. Heije, D. Valerio and B.J.E.G. Bast (The Netherlands) 1

Simultaneous detection of allotypes in native and activated human C2 by isoelectric focusing and silver staining (JIM 05045)
O.G. Segurado and A. Arnaiz-Villena (Spain) 9

A new graphical method for determining the affinity constants of monoclonal antibodies to enzymes (JIM 05046)
H. Ehle, C. Gödicke and A. Horn (G.D.R.) 17

Determination of epitope specificities and affinities of monoclonal antibodies in solution phase using biotin-labeled carcinoembryonic antigen and avidin as precipitating agent (JIM 05047)
U. Krüger, L. Wickert and C. Wagener (F.R.G.) 25

Time-resolved fluoroimmunoassay of 5-methyl-2′-deoxycytidine employing europium-labeled antigen as tracer (JIM 05048)
S. Rasi, E. Suvanto, L.M. Vilpo and J.A. Vilpo (Finland) 33

A specific solid-phase assay for the detection of immune complexes containing *Entamoeba histolytica* antigens (JIM 05049)
A. Mohimen, T.K. Maitra, K.N. Jalan and S. Mehra (India) 39

Multiple immunoenzyme staining techniques. Use of fluoresceinated, biotinylated and unlabelled monoclonal antibodies (JIM 05050)
C.M. Van der Loos, P.K. Das, J.J. Van den Oord and H.J. Houthoff (The Netherlands) 45

Simultaneous measurement by flow cytometry of phagocytosis and hydrogen peroxide production of neutrophils in whole blood (JIM 05051)
M. Hasui, Y. Hirabayashi and Y. Kobayashi (Japan) 53

Enzyme immunoassay detection of circulating immune complexes by monoclonal antibodies to C3 neoepitopes with special reference to IgG concentration and to interfering anti-immunoglobulin antibodies (JIM 05053)
P. Garred, T.E. Mollnes, T. Lea and P.J. Lachmann (Norway, U.K.) 59

Development of polyclonal and monoclonal antibodies for immunoassay and neutralization of human interleukin-4 (JIM 05054)
I. Chrétien, A. Van Kimmenade, M.K. Pearce, J. Banchereau and J.S. Abrams (France, U.S.A.) 67

Detection and reduction of Protein A contamination in immobilized Protein A purified monoclonal antibody preparations (JIM 05055)
J.W. Bloom, M.F. Wong and G. Mitra (U.S.A.) 83

Comparison of antifading agents used in immunofluorescence (JIM 05056)
K.D. Krenik, G.M. Kephart, K.P. Offord, S.L. Dunnette and G.J. Gleich (U.S.A.) 91

Rapid screening and selection of monoclonal antibodies by bivariate flow cytometric analyses (JIM 05057)
M.G. Pallavicini, S. Haendel, C. Rosette and R. Barlett (U.S.A.) 99

A biotin-avidin sandwich ELISA for quantification of intact complement component C9. The sera from hereditary C9 deficient individuals completely lack C9 (JIM 05058)
Y. Takata, T. Moriyama, Y. Fukumori, A. Yoden, M. Shima and S. Inai (Japan) 107

Quantitation of a cationic antimicrobial granule protein of human polymorphonuclear leukocytes by ELISA (JIM 05059)
H.A. Pereira, L.E. Martin and J.K. Spitznagel (U.S.A.) 115

Production and characterization of monoclonal antibodies against human glycoalbumin (JIM 05060)
M.P. Cohen and E. Hud (U.S.A.) 121

Purification of monoclonal antibodies raised against prostate-specific acid phosphatase for use in vivo in radioimaging of prostatic cancer (JIM 05062)
L. Hakalahti and P. Vihko (Finland) 131

Short communications

An improved latex agglutination-inhibition test for the detection of human chorionic gonadotropin in urine (JIM 05088)
P.-L. Lim and W.-F. Choy (Hong Kong) 137

A blocking agent and a blocking step are not needed in ELISA, immunostaining dot-blots and Western blots (JIM 05092)
K. Mohammad and A. Esen (U.S.A.) 141

0022-1759/89/$03.50 © 1989 Elsevier Science Publishers B.V. (Biomedical Division)

Journal of Immunological Methods, 117 (1989) 296–297
Elsevier

Contents to volume 117

Number 1 – 8 February 1989

Research reports

Production of hybrid hybridomas based on HAT^s-neomycin^r double mutants (JIM 05043)
W.B.M. De Lau, A.E. Van Loon, K. Heije, D. Valerio and B.J.E.G. Bast (The Netherlands) 1

Simultaneous detection of allotypes in native and activated human C2 by isoelectric focusing and silver staining (JIM 05045)
O.G. Segurado and A. Arnaiz-Villena (Spain) 9

A new graphical method for determining the affinity constants of monoclonal antibodies to enzymes (JIM 05046)
H. Ehle, C. Gödicke and A. Horn (G.D.R.) 17

Determination of epitope specificities and affinities of monoclonal antibodies in solution phase using biotin-labeled carcinoembryonic antigen and avidin as precipitating agent (JIM 05047)
U. Krüger, L. Wickert and C. Wagener (F.R.G.) 25

Time-resolved fluoroimmunoassay of 5-methyl-2′-deoxycytidine employing europium-labeled antigen as tracer (JIM 05048)
S. Rasi, E. Suvanto, L.M. Vilpo and J.A. Vilpo (Finland) 33

A specific solid-phase assay for the detection of immune complexes containing *Entamoeba histolytica* antigens (JIM 05049)
A. Mohimen, T.K. Maitra, K.N. Jalan and S. Mehra (India) 39

Multiple immunoenzyme staining techniques. Use of fluoresceinated, biotinylated and unlabelled monoclonal antibodies (JIM 05050)
C.M. Van der Loos, P.K. Das, J.J. Van den Oord and H.J. Houthoff (The Netherlands) 45

Simultaneous measurement by flow cytometry of phagocytosis and hydrogen peroxide production of neutrophils in whole blood (JIM 05051)
M. Hasui, Y. Hirabayashi and Y. Kobayashi (Japan) 53

Enzyme immunoassay detection of circulating immune complexes by monoclonal antibodies to C3 neoepitopes with special reference to IgG concentration and to interfering anti-immunoglobulin antibodies (JIM 05053)
P. Garred, T.E. Mollnes, T. Lea and P.J. Lachmann (Norway, U.K.) 59

Development of polyclonal and monoclonal antibodies for immunoassay and neutralization of human interleukin-4 (JIM 05054)
I. Chrétien, A. Van Kimmenade, M.K. Pearce, J. Banchereau and J.S. Abrams (France, U.S.A.) 67

Detection and reduction of Protein A contamination in immobilized Protein A purified monoclonal antibody preparations (JIM 05055)
J.W. Bloom, M.F. Wong and G. Mitra (U.S.A.) 83

Comparison of antifading agents used in immunofluorescence (JIM 05056)
K.D. Krenik, G.M. Kephart, K.P. Offord, S.L. Dunnette and G.J. Gleich (U.S.A.) 91

Rapid screening and selection of monoclonal antibodies by bivariate flow cytometric analyses (JIM 05057)
M.G. Pallavicini, S. Haendel, C. Rosette and R. Barlett (U.S.A.) 99

A biotin-avidin sandwich ELISA for quantification of intact complement component C9. The sera from hereditary C9 deficient individuals completely lack C9 (JIM 05058)
Y. Takata, T. Moriyama, Y. Fukumori, A. Yoden, M. Shima and S. Inai (Japan) 107

Quantitation of a cationic antimicrobial granule protein of human polymorphonuclear leukocytes by ELISA (JIM 05059)
H.A. Pereira, L.E. Martin and J.K. Spitznagel (U.S.A.) 115

Production and characterization of monoclonal antibodies against human glycoalbumin (JIM 05060)
M.P. Cohen and E. Hud (U.S.A.) 121

Purification of monoclonal antibodies raised against prostate-specific acid phosphatase for use in vivo in radioimaging of prostatic cancer (JIM 05062)
L. Hakalahti and P. Vihko (Finland) 131

Short communications

An improved latex agglutination-inhibition test for the detection of human chorionic gonadotropin in urine (JIM 05088)
P.-L. Lim and W.-F. Choy (Hong Kong) 137

A blocking agent and a blocking step are not needed in ELISA, immunostaining dot-blots and Western blots (JIM 05092)
K. Mohammad and A. Esen (U.S.A.) 141

0022-1759/89/$03.50 © 1989 Elsevier Science Publishers B.V. (Biomedical Division)

Corrigendum
Biotinylated recombinant interleukin-2. A tool for research on the interleukin-2 receptor (JIM 05101)
B.M.J. Foxwell, D. Taylor, B. Greiner, M.J. Mihatsch, V. Olivieri and B. Ryffel (Switzerland) 147

Book reviews 149

Announcement 151

Notes to authors 152

Number 2 – 24 February 1989

Research reports
Rapid assay and identification of human haemopoietic growth factors (JIM 05063)
W.E. Pullman, S. Elsbury, M. Kobayashi, J. Hargreaves, L. Preston, B. Van Leeuwen and A.J. Hapel (Australia) 153
Purification of human basophils. Their response to anti-IgE (JIM 05064)
M.A. Lett-Brown, L. Robinson, H.S. Juneja and J.A. Grant (U.S.A.) 163
Murine monoclonal anti-avidin antibodies enhance the sensitivity of avidin-biotin immunoassays and immunohistologic staining (JIM 05065)
W.F. Cassano (U.S.A.) 169
Quantitative assay of a human monoclonal IgM antibody (HA-1A) in human serum (JIM 05066)
M.B. Khazaeli, K.J. Rogers and A.F. LoBuglio (U.S.A.) 175
Kinetic assay of cytotoxic T lymphocyte clones. I. Methodology and data analysis (JIM 05067)
L.J. Jenski and R.M. Kleyle (U.S.A.) 181
Kinetic assay of cytotoxic T lymphocyte clones. II. Comparison of primary and secondary responses and lectin-dependent cytotoxicity (JIM 05068)
L.J. Jenski (U.S.A.) 191
ELISA-plaque assay for the detection of platelet antibody-secreting cells. I. Studies on optimization of reagent concentrations and culture conditions for enumeration of spot-forming cells (JIM 05069)
H. Uchiyama and J. Akatsuka (Japan) 199
Cell-mediated cytotoxicity. A highly sensitive and informative flow cytometric assay (JIM 05070)
S.E. Slezak and P.K. Horan (U.S.A.) 205
Cross-reaction of antibodies to coupling groups used in the production of anti-peptide antibodies (JIM 05073)
R.J. Edwards, A.M. Singleton, A.R. Boobis and D.S. Davies (U.K.) 215
Specificity of two-site immunoassays (JIM 05074)
L.M. Boscato, G.M. Egan and M.C. Stuart (Australia) 221
Partial purification and characterization of a factor produced by mitogen-stimulated sheep mononuclear cells with thymocyte-activating and cartilage-degrading properties (JIM 05075)
G.D. Harkiss, K. Price and D. Veitch (U.K.) 231
A method for the quantitation of interleukin-2 activity (JIM 05076)
G. Hewlett, K.G. Stünkel and H.D. Schlumberger (F.R.G.) 243
Assays for total and antigen-specific polymeric IgA in serum based on binding to secretory component (JIM 05093)
R.C.A. Bartholomeusz, J.T. LaBrooy, P.L. Ey, A.C. Di Matteo, D.A. Daniels, C.S. Anderson and D. Rowley (Australia) 247
Solid-phase immunoassay of serum antibodies to peptides. Covalent antigen binding to adsorbed phenylalanine-lysine copolymers (JIM 05095)
R.N. Hobbs (U.K.) 257
A two-particle turbidometric latex immunoassay for the detection of specific IgM antibodies (JIM 05097)
P.-L. Lim, K.-H. Ko and W.-F. Choy (Hong Kong) 267
Detection and isolation of antigen-specific B cells by the fluorescence activated cell sorter (FACS) (JIM 05123)
M.Y. Hoven, L. De Leij, J.F.K. Keij and T.H. The (The Netherlands) 275

Short communications
Analysis of *Candida albicans* allergen extracts with a novel crossed enzyme immunoelectrophoresis (CEIE) technique. Binding of the colour precipitate to a nitrocellulose membrane (JIM 05117)
J. Savolainen and M. Viander (Finland) 285
Separation of B and T lymphocytes by a hybrid field-flow fractionation/adhesion chromatography technique (JIM 05119)
J.C. Bigelow, J.C. Giddings, Y. Nabeshima, T. Tsuruta, K. Kataoka, T. Okano, N. Yui and Y. Sakurai (U.S.A., Japan) 289

Book review 295

Contents to volume 117 (1989) 296

Notes to authors 298